CUMULATIVE SUBJECT INDEX

Environmental Entomology,

Volume 15, 1986

Abundance, *Oryctes rhinoceros*, 84
Acalymma vittatum, *Aphis gossypii*, 963
Diabrotica balteata, 963
Diabrotica undecimpunctata howardi, 963
dispersion indices, 963
Empoasca abrupta, 963
Acleris minuta, developmental polymorphism, 258
ericaceous plants, 258
rosaceous plants, 258
Acoloithus falsarius, attractant, 959
Harrisina americana, 959
pheromone, 959
Acyrthosiphon kondoi, development, 227
phenology models, 227
population, 91
Acyrthosiphon pisum, control by *Verticillium lecanii*, 281
development, 227
phenology model, 227
Adelges cooleyi, population, 254
sampling, 254
seasonal abundance, 254
spatial abundance, 254
Adelges picese, Fraser fir, 911
water stress, 911
Adult diapause, *Beauveria bassiana*, 884
Conotrachelus nenuphar, 884
migration, 884
Aedes atlanticus, *Aedes vexans*, 874
body size, 874
Culex salinarius, 874
feeding success, 874
Psorophora columbiae, 874
Aedes vexans, *Aedes atlanticus*, 874
body size, 874
Culex salinarius, 874
feeding success, 874
Psorophora columbiae, 874
Aflatoxin, *Aspergillus flavus*, 1116
Heliothis zea, 1116
maize husk, 1116
Ostrinia nubilalis, 1116
Aggregation, *Hylobius abietis*, 1050
host odors, 1050
Agroecosystem diversification, arthropod natural enemies, 456
Agromyza frontella, biological control, 448
Chrysocharis punctifacies, 448
Dacnusa dryas, 448
Miscogaster hortensis, 448
Agrotis ipsilon, *Bonnetia comta*, 360
nuclear polyhedrosis virus, 360
Rachiplusia ou, 360
Agrotis orthogonia, chlorpyrifos, 147
horned larks, 147
McCown's longspurs, 147
population, 147
wheat, 147
Agrypon flaveolatum, biological control, 1042
Cysenis albicans, 1042
Operophtera brumata, 1042
parasitism, 1042
Agrytaenia citrana, capture of males, 232
pheromone traps, 232
virgin females, 232
Alabama argillacea, cotton leaves, 128
development, 128
feeding, 128
glandular chemicals, 128
survival, 128
Aldrin, *Diabrotica barberi*, 427
geographic differences, 427
Alfalfa, glandular-haired versus nonglandular, 396
Lygus desertinus, 132
Lygus elisus, 132
Lygus hesperus, 132
resistance to *Hypera postica*, 396
Thrips tabaci, 695
Alfalfa seeds, behavior, 1244
Bruchophagus roddi, 388
chalcids, 1244
morphological changes, 388
seed development, 388
volatile components, 1244
Almond leaves, damage, 190
gas exchange rates, 594
Panonychus citri, 190
Panonychus ulmi, 190
spider mites, 594
Tetranychus pacificus, 190
Tetranychus urticae, 190
water potential, 594
water stress, 594
Alsophila pometaria, behavior, 943
communication disruption, 943
pheromone, 943
Amblyseius californicus, biological control, 790
corn, 790
Oligonychus pratensis, 790
Phytoseilus persimilis, 790
Tetranychus urticae, 790
Anagasta kuehniella, male response, 369
pheromone communication, 369
temporal differences, 369
wind tunnel, 369
Anaphes sordidatus, biology, 100
life cycle, 100
Listronotus oregonensis 100
on *Listronotus oregonensis*, 469
parasitization, 469
Anasa tristis, in polycultures, 285
Antibiosis tests, design of experiments, 581
optimum sample size, 581
Schizaphis graminum, 581
Anticarsia gemmatalis, color phase, 784
density, 779, 784
Geocoris punctipes, 1032
host-plant phenology, 784
host-plant resistance, 1032
larval development, 784
Mocis disseverans, 1272
nuclear polyhedrosis virus, 573
Pseudoplusia includens, 1032
population dynamics, 809
predators, 1032
sex pheromone, 1272
soybeans, 573, 779, 809, 1032
Aonidiella aurantii, light intensity, 971
male emergence, 971
relative humidity, 971
temperature, 971
Apanteles fumiferanae, *Choristoneura pinus*, 409
Choristoneura pinus, 662
hyperparasitism, 662
Michigan, 409
mortality, 662
parasitization, 662
spatial activity, 409
temporal activity, 409
Apanteles morrisi, *Choristoneura pinus*, 409
Choristoneura pinus, 662
hyperparasitism, 662
Michigan, 409
mortality, 662
parasitization, 662
spatial activity, 409
temporal activity, 409
Aphid eggs, *Brachycolus asparagi*, 989
Brachycorynella asparagi, 989
desiccation, 989
humidity, 989
Aphidius lupini, *Macrosiphum albifrons*, 719
Aphidoletes aphidimyza, diapause, 1067
low-intensity radiation, 1067
Aphids, discriminant analysis, 1145
parasitoids, 1171
plant susceptibility, 1145
shade trees, 1145
urban entomology, 1145
Aphis gossypii, *Acalymma vittatum*, 963
Diabrotica balteata, 963
Diabrotica undecimpunctata howardi, 963
dispersion indices, 963
Empoasca abrupta, 963
Aphis pomi, age, 614
developmental rate, 607,614
ovarian parameters, 607
parental rearing conditions, 614
reproduction, 614
temperature effects, 607
weight, 607, 614
Apis mellifera, apples, 723, 1047
B exotoxin, 166
Bacillus thuringiensis, 166
cross-pollination, 723
fruit set, 723
fungicides, 1047
in-hive pollination, 723
insecticides, 170
knockdown, 170
methyl parathion, 170
mortality, 166, 170
nestmate contact, 723
pears, 1047
Penncap-M, 170
thuringiensin, 166
toxicity, 166, 170
Apples, *Apis mellifera*, 1047
Apis mellifera, 723
developmental thresholds, 984
fungicides, 1047
Orius insidiosus, 984
Phyllonrycter elmaella, 635
predation, 984
Araneae, ballooning, 1225
dispersal, 1225
male behavior, 1225
Arrested larval development, crowding, 1264
overfeeding, 1264
Tribolium castaneum, 1264

Arthropod natural enemies, agroecosystem diversification
effects, 456
behavior, 456
biological control, 456
Arthropods, collards, 826
sewage sludge, 826
Artificial diet, *Spodoptera frugiperda*, 327
Artificial seeds, *Callosobruchus maculatus*, 1286
cowpeas, 1286
growth inhibitors, 1286
Artogeia rapae, host/parasite effects, 1234
larval density, 1234
Pteromalus puparum, 994
parasitism, 994
Aspergillus flavus, aflatoxin, 1116
Heliothis zea, 1116
maize husk, 1116
Ostrinia nubilalis, 1116
Associates, exclusion, 1104
Ips calligraphus, 1104
mortality, 1104
Attagenus unicolor, pupation, 1166
survival, 1166
Attractant, *Acoloithus falsarius*, 959
Harrisina americana, 959
pheromone, 959

B exotoxin, *Bacillus thuringiensis*, 166
thuringiensin, 166
toxicity, 166
Bacillus popilliae, *Cyclocephala parallela*, 703
Ligyrus subtropicus, 703
sugarcane, 703
Bacillus thuringiensis, *Apis mellifera*, 166
B exotoxin, 166
bioassay, 1161
Heliothis virescens, 1161
Heliothis zea, 1161
nuclear polyhedrosis virus, 1161
thuringiensin, 166
Ballooning, Araneae, 1225
dispersal, 1225
male behavior, 1225
Beans, *Evarthrus sodalis*, 1028
Harpalus pennsylvanicus, 1028
intercropping, 1028
Beauveria bassiana, adult diapause, 884
Cerotoma sp., 642
Chalcodermus aeneus, 642
Chrysomelidae, 1237
Conotrachelus nenuphar, 884
conidia survival, 642
Metarhizium anisopliae, 1237
migration, 884
sunlight, 1237
Behavior, agroecosystem diversification, 456
alfalfa seed, 1244
Alsophila pometaria, 943
arthropod natural enemies, 456
Blatella germanica, 198
Bruchophagus roddi, 388
chalcids, 1244
communication disruption, 943
Heliothis phloxiphaga, 931
pheromone identification, 931
pheromone, 943
sex pheromone, 931
volatile components, 1244
Bell pepper, infestation by *Liriomyza trifolii*, 383
Liriomyza trifolii, 96
Bemisia tabaci, CO_2, 61
cotton, 61
developmental rate, 1179
in cotton, 300
in lettuce, 300
parasites, 1179
plant viruses, 1179
population, 61
Trialeurodes abutilonea, 300
trap, 300
weed hosts, 1179
Beneficial insects, *Ips* spp., 1220
pine trees, 1220
Bermudagrass, development, 889
Spodoptera frugiperda, 889
Bessa harveyi, life table, 246
mortality, 246
parasite of *Pikonema alaskensis*, 246
Betulaphis brevipilosa, European white birch, 771
tree species determination, 771
Bexotoxin, *Apis mellifera*, 166
Bindweed, host range, 834
Megacerus discoidus, 834
Bioassay, *Bacillus thuringiensis* 1161
Heliothis virescens, 1161
Heliothis zea, 1161
nuclear polyhedrosis virus, 1161
Biological control, *Agromyza frontella*, 448
Agrypon flavicolatum, 1042
Amblyseius californicus, 790
arthropod natural enemies, 456
Cyclosa turbinata, 177
Cyzenis albicans, 1042
corn, 790
Dialeurodes citri, 585
development rate, 1301
Encarsia lahorensis, 585
fungi, 1070
house flies, 1296
insecticide resistance, 1255
Metepeira grinnelli, 177
Musca domestica, 436
Muscidifurax zaraptor, 1296
Nasonia vitripennis, 436
Nilaparvata lugens, 1070
Oligonychus pratensis, 790
Operophtera brumata, 1042
Phytoseilus persimilis, 790
parasites, 1268
parasitism, 1042
pesticide-resistant natural enemies, 1255
phytoseiid mites, 1255
Rhopalomyia californica, 1268
rice, 1070
Spalangia cameroni, 1296
stable flies, 1296
survivorship, 1301
Tetranychus urticae, 790
Tetrastichus sp., 1268
Torymus baccharidis, 1268
Urolepis rufipes, 1296, 1301
Verticillium lecanii, 380
Biology, density-related stress, 914
feeding rhythms, 914
Lymantria dispar, 914
larval development, 914
naphes sordidatus 100
Biotypic status, *Schizaphis graminum*, 567
Blattella germanica, behavior, 198
density, 198
drinking, 198
feeding, 198
harborage-to-resource distance, 198
mortality, 221
yeast infection, 221
Bluestain, *Ceratocystis minor*, 951
Dendroctonus frontalis, 951
Tarsonemus ips, 951
Tarsonemus krantzi, 951
Body size, *Aedes atlanticus*, 874
Aedes vexans, 874
Culex salinarius, 874
feeding success, 874
Psorophora columbiae, 874
Boisea rubrolineatus, forest structure, 1055
population dynamics, 1055
Boll weevil, *Bracon mellitor*, 880
host insect/plant relationships, 880
Bombus flavifrons, flower availability, 750
reproduction, 750
temperature, 750
Bonnetia comta, *Agrotis ipsilon*, 360
corn, 360
nuclear polyhedrosis virus, 360
Rachiplusia ou, 360
Boundary layer, Cicadellidae, 1199
migration, 1199
oogenesis/flight syndrome, 1199
Brachycolus asparagi, aphid eggs, 989
Brachycorynella asparagi, 989
desiccation, 989
humidity, 989
Brachycorynella asparagi, aphid eggs, 989
Brachycolus asparagi, 989
desiccation, 989
humidity, 989
Bracon hebetor, *Cadra cautella*, 767
natural enemies, 767
Plodia interpunctella, 767
peanuts, 767
Bracon mellitor, boll weevil, 880
host insect/plant relationships, 880
Breeding, *Oryctes rhinoceros*, 84
Psorophora ferox, 683
Brevicoryne brassicae, on cabbage in bare-ground monocultures, 293
on cabbage in living mulches, 293
Bruchophagus roddi, behavior, 388
oviposition, 388
seed development, 388
seed morphological changes, 388
site selection, 388
Burlap bands, effect on between-tree movement, 373
Lymantria dispar, 373
monitoring tool, 373

Cabbage, *Thrips tabaci*, 513, 520
Cadra cautella, *Bracon hebetor*, 767
male response, 369
natural enemies, 767
peanuts, 767
pheromone communication, 369
temporal differences, 369
wind tunnel, 369
Xylocoris flavipes, 767
Calaphis flava, European white birch, 771
tree species determination, 771
Callipterinella calliptera, European white birch, 771
tree species determination, 771
Callosobruchus maculatus, artificial seeds, 1286
cowpeas, 1286
growth inhibitors, 1286
Carabids, corn agroecosystem, 1017
predation, 1017
predator activity, 1017
Carbaryl, *Chrysoperla carnea*, 1130
pesticides, 1130
resistance, 1130
Ceratitis capitata, annual occurrence, 507
distance/response curves, 71
host fruits, 507
mortality, 465
population, 507
seasonal occurrence, 507
Steinernema feltiae, 465
trapping, 71

traps, 507
trimedlure, 71
Ceratocystis, disturbance, 1090
epidemiology, 1090
forests, 1090
Ceratocystis minor, bluestain, 951
Dendroctonus frontalis, 951
Tarsonemus ips, 951
Tarsonemus krantzi, 951
Cerotoma sp., *Beauveria bassiana*, 642
Chaetocnema pulicaria, sweet corn, 867
traps, 867
Chalcids, alfalfa seed, 1244
behavior, 1244
volatile components, 1244
Chalcodermus aeneus, *Beauveria bassiana*, 642
Chemical exclusion, filberts, 1037
Myzocallis coryli, 1037
predators, 1037
Chemo-orientation, *Diabrotica virgifera virgifera*, 839
host location, 839
olfaction, 839
Cheumatopsyche pettiti, behavior, 355
flight patterns, 355
swarming, 355
3-D analysis, 355
Chilo partellus, *Eldana saccharina*, 1124
maize, 1124
resistance, 1124
Chilo suppressalis, development, 417
on modern rice cultivars, 417
on traditional rice cultivars, 417
population, 417
Chilomenes sexmaculata, development period, 800
fecundity, 800
food consumption, 800
longevity, 800
temperature, 800
Chlorpyrifos, *Agrotis orthogonia*, 147
horned larks, 147
McCown's longspurs, 147
Choristoneura fumiferana, monitoring, 152
pheromones, 152
population, 152
traps, 152
Choristoneura occidentalis, distribution, 547
stochastic phenology model, 547
Choristoneura pinus, *Apanteles fumiferanae*, 662
Apanteles morrisi, 662
Chromaphis juglandicola, microhabitat distribution, 555
spatial dispersion, 555
Chrysanthemums, *Liriomyza trifolii*, 268
oviposition behavior, 1189
Spodoptera exigua, 1189
Chrysocharis punctifacies, *Agromyza frontella*, 448
biological control, 448
Chrysomelidae, *Beauveria bassiana*, 1237
Metarhizium anisopliae, 1237
sunlight, 1237
Chrysoperla carnea, carbaryl, 1124
pesticides, 1130
resistance, 1130
Cicadellidae, boundary layer, 1199
migration, 1199
oogenesis/flight syndrome, 1199
Circadian pattern, cow color, 1307
Haematobia irritans, 1307
Cnaphalocrocis medinalis, development, 417
on modern rice cultivars, 417
on traditional rice cultivars, 417
population, 417
Cochylis hospes, sex pheromones, 1063
Coconuts, *Oryctes rhinoceros*, 84
Cold acclimation, diapause, 936
Muscidifurax zaraptor, 936
Spalangia cameroni, 936
Urolepis rufipes, 936
Collards, arthropods, 826
sewage sludge, 826
Colonization, *Leptinotarsa decemlineata*, 311
Color phase, *Anticarsia gemmatalis*, 784
density, 784
host-plant phenology, 784
larval development, 784
Colored sticky panels, Hymenoptera, 64
Communication disruption, *Alsophila pometaria*, 943
behavior, 943
pheromone, 943
Compensatory processes, *Manduca sexta*, 1137
Nicotiana tabacum, 1137
simulated defoliation, 1137
yield loss, 1137
Competition, *Conopthorus resinosae*, 88
Dioryctria disclusa, 88
Computer program, degree-day, 999
physiological time, 999
thermal unit, 999
Conidia survival, *Beauveria bassiana*, 642
Conidial germination, *Delia antiqua*, 1154
Entomophthora muscae, 1154
moisture, 1154
temperature, 1154
Conopthorus resinosae, competition, 88
Dioryctria disdusa, 88
feeding, 88
red pine, 88
Conotrachelus senuphar, adult diapause, 884
Beauveria bassiana, 884
migration, 884
Corn, egg-mass detrition, 817
Graminella nigrifrons, 1119
Johnsongrass, 1119
life tables, 1119
Oligonychus pratensis, 79
Ostrinia nubilalis, 106, 501, 631, 817
Sitophilus zeamais, 654
corn agroecosystem, carabids, 1017
predation, 1017
predator activity, 1017
Cotton, *Bemisia tabaci*, 61
Crocidosema plebejana, 1078
damage, 399
Frankliniella occidentalis, 821
functional response, 821
Heliothis zea, 365
Lygus desertinus, 132
Lygus elisus, 132
Lygus hesperus, 132
Ostrinia nubilalis, 399
omnivore, 821
opportunist, 821
predator, 821
spider mites, 821
Trichogramma pretiosum, 365
trichome density, 365
weather, 1078
Cotton leaves, *Alabama argillacea*, 128
Cow color, circadian pattern, 1307
Haematobia irritans, 1307
Cowpeas, artificial seeds, 1286
Callosobruchus maculatus, 1286
growth inhibitors, 1286
Crocidosema plebejana, cotton, 1078
weather, 1078
Cross-pollination, *Apis mellifera*, 723
Crowding, arrested larval development, 1264
overfeeding, 1264
Sitophilus zeamais, 733
Tribolium castaneum, 1264
Culex salinarius, *Aedes atlanticus*, 874
Aedes vexans, 874
body size, 874
feeding success, 874
Psorophora columbiae, 874
Cyclocephala paralilela, *Bacillus popilliae*, 703
sugarcane, 703
Ligyrus subtropicus, 703
Cyclocephala pasadenae, flight, 194
Phyllophaga crinita, 194
traps, 194
Cyclosa turbinata, biological control, 177
interspecific competition, 177
Metepeira grinnelli, 177
on insect populations, 177
predators, 177
Cysenis albicans, *Agrypon flaveolatum*, 1042
biological control, 1042
Operophtera brumata, 1042
parasitism, 1042

Dacnusa dryas, *Agromyza frontella*, 448
biological control, 448
Dacus cucurbitae, distribution, 488
host plants, 488
mortality, 465
Steinernema feltiae, 465
Dacus dorsalis, mortality, 465
Steinernema feltiae, 465
Degree-day, computer program, 999
physiological time, 999
thermal unit, 999
Delia antiqua, conidial germination, 1154
Entomophthora muscae, 1154
moisture, 1154
temperature, 1154
Delia florilega, *Delia platura*, 1022
development rates, 1022
Delia platura, *Delia florilega*, 1022
development rates, 1022
Dendroctonus frontalis, bluestain, 951
Ceratocystis minor, 951
Tarsonemus ips, 951
Tarsonemus krantzi, 951
Dendroctonus spp., disturbance, 850
guilds, 850
Ips spp., 850
population dynamics, 850
Dendroctonus terebrans, in-flight responses, 710
monoterpene attraction, 710
turpentine attraction, 710
Density, *Anticarsia gemmatalis*, 779
Anticarsia gemmatalis, 784
color phase, 784
host-plant phenology, 784
larval development, 784
Oligonychus pratensis, 79
soybeans, 779
Density relationships, *Laplatacris dispar*, 775
mortality, 775
population dynamics, 775
Density/height profile, *Lymantria dispar*, 431
Density-related stress, biology, 914
feeding rhythms, 914
Lymantria dispar, 914
larval development, 914
Desiccation, aphid eggs, 989
Brachycolus asparagi, 989
Brachycorynella asparagi, 989
humidity, 989
Development, *Acleris minuta*, 258
Acyrthosiphon kondoi, 227
Acyrthosiphon pisum, 227
Alabama argillacea, 128
Aphis pomi, 607, 614
bermudagrass, 889
Chilo suppressalis, 417
Cnaphalocrocis medinalis, 417

Diabrotica undecimpunctata howardi, 626
Diabrotica virgifera virgifera, 626
forage crops, 889
Monochamus carolinensis, 251
Nabis roseipennis, 536
Nephotettix virescens, 417
Nezara viridula, 739
Nilaparvata lugens, 417
Nymphula depunctalis, 417
Platynota idaeusalis, 210
Sogatella furcifera, 417
Spodoptera frugiperda, 182, 889
Development period, *Chilomenes sexmaculata*, 800
fecundity, 800
food consumption, 800
longevity, 800
temperature, 800
Development rate, biological control, 1301
survivorship, 1301
Urolepis rufipes, 1301
Development rates, *Delia florilega*, 1022
Delia platura, 1022
Developmental rate, *Semiscia tabaci*, 1179
parasites, 1179
plant viruses, 1179
weed hosts, 1179
Developmental thresholds, apples, 984
Orius insidiosus, 984
predation, 984
Diabrotica balteata, *Acalymma vittatum*, 963
Aphis gossypii, 963
Diabrotica undecimpunctata howardi, 963
dispersion indices, 963
Empoasca abrupta, 963
Diabrotica barberi, aldrin susceptibility, 427
population, 427
Diabrotica speciosa, *Beauveria bassiana*, 642
Diabrotica undecimpunctata howardi, *Acalymma vittatum*, 963
Aphis gossypii, 963
Diabrotica balteata, 963
Diabrotica virgifera virgifera, 626
development, 626
dispersion indices, 863
Empoasca abrupta, 963
humidity effects, 626
survival, 626
temperature effects, 626
Diabrotica virgifera virgifera, chemo-orientation, 839
development, 626
Diabrotica undecimpunctata howardi, 626
heat units, 109
host location, 839
humidity effects, 626
migratory flight, 620
olfaction, 839
phenology, 109
planting dates, 109
population, 109
survival, 626
temperature effects, 626
Dialeurodes citri, biological control, 585
Encarsia lahorensis, 585
population, 585
Diapause, cold acclimation, 936
commitment, 669
development of *Papaipema nebris* eggs, 403
Haematobia irritans, 44
Heliothis virescens, 919
Heliothis zea, 919
induction, 669
Musca autumnalis, 669
Muscidifurax zaraptor, 936
Pectinophora gossypiella, 494
Spalangia cameroni, 936
thermal effects, 919
Urolepis rufipes, 936
Diaphania hyalinata, polycultures, 285
population, 285
squash, 285
Diel activity, *Musca domestica*, 56
Diet, *Nabis roseipennis*, 536
Dioryctria amatella, pine, 316
sex pheromone, 316
trap, 316
Dioryctria disclusa, competition, 88
Conopthorus resinosae, 88
feeding, 88
red pine, 88
Discriminant analysis, aphids, 1145
plant susceptibility, 1145
shade trees, 1145
urban entomology, 1145
Dispersal, Araneae, 1225
ballooning, 1225
Lymantria dispar, 431
male behavior, 1225
Dispersion indices, *Acalymma vittatum*, 963
Aphis gossypii, 963
Diabrotica balteata, 963
Diabrotica undecimpunctata howardi, 963
Empoasca abrupta, 963
Dispersion pattern, *Liriomyza trifolii*, 383
Dispersion patterns, *Oligonychus pratensis*, 335
Tetranychus urticae, 335
Distribution, *Choristoneura occidentalis*, 547
Dacus cucurbitae, 488
Liriomyza trifolii, 96
Ostrinia nubilalis, 106, 399
Disturbance, *Ceratocystis*, 1090
Dendroctonus spp., 850
epidemiology, 1090
forests, 1090
guilds, 850
Ips spp., 850
population dynamics, 850
Drosophila melanogaster, *Drosophila simulans*, 859
Leptopilina boulardi, 859
seasonal development, 859
synchrony, 859
Drosophila simulans, *Drosophila melanogaster*, 859
Leptopilina boulardi, 859
seasonal development, 859
synchrony, 859
Drought stress, *Schizaphis graminum*, 118

Egg melanization, *Sitona lineatus*, 601
Egg parasites, longevity, 1149
Lymantria dispar, 1149
Ooencyrtus kuvanae, 1149
photoperiod, 1149
progeny production, 1149
Egg-mass detrition, corn, 817
Ostrinia nubilalis, 817
Egg-mass dispersion, *Ostrinia nubilalis*, 1212
sampling, 1212
Eggs, *Psorophora ferox*, 683
Elasmopalpus lignosellus, grain sorghum, 905
larval distribution, 905
larval phenology, 905
longevity, 715
oviposition rate, 715
temperatures, 715
Eldana saccharina, *Chilo partellus*, 1124
maize, 1124
resistance, 1124
Elm, host suitability, 843
Xanthogaleruca luteola, 143, 843
Empoasca abrupta, *Acalymma vittatum*, 963
Aphis gossypii, 963
Diabrotica balteata, 963
Diabrotica undecimpunctata howardi, 963
dispersion indices, 963
Empoasca fabae, population dynamics, 898
Encarsia lahorensis, biological control, 585
Dialeurodes citri, 585
gardenias, 585
Endophytic fungus, host preference, 1083
Spodoptera frugiperda, 1083
tall fescue, 1083
Energy budget, *Malacosoma americanum*, 795
nutritional ecology, 795
Entomophthora muscae, conidial germination, 1154
Delia antiqua, 1154
moisture, 1154
temperature, 1154
Entomophthora sp., *Macrosiphum albifrons*, 719
Eoreuma loftini, ovipositional sites, 75
parasitized by Trichogramma
retorridum, 75
sugarcane, 75
Ephedrus californicus, *Macrosiphum albifrons*, 719
Epidemiology, *Ceratocystis*, 1090
disturbance, 1090
forests, 1090
Epilachna varivestis, fluazifop-butyl, 216
growth-regulator herbicides, 216
mefluidide, 216
microspondia, 32
mortality, 32
Nosema epilachnae, 32
Nosema varivestis, 32
on lima bean, 216
on soybean, 216
Pediobius foveolatus host, 32
sethoxydim, 216
Eucelatoria bryani, *Heliothis* spp., 1111
parasitoids, 1111
proteins, 1111
Euceraphis betulae, European white birch, 771
tree species determination, 771
Eumegachile pugnata, mortality, 877
nesting, 877
temperature, 877
European white birch, aphid host preference, 771
Betulaphis brevipilosa, 771
Calaphis flava, 771
Callipterinella callipterа, 771
Euceraphis betulae, 771
species determination, 771
Evarthrus sodalis, beans, 1028
Harpalus pennsylvanicus, 1028
intercropping, 1028
tomatoes, 1028
Exclusion, associates, 1104
Ips calligraphus, 1104
mortality, 1104

Fecundity, *Chilomenes sexmaculata*, 800
development period, 800
food consumption, 800
life tables, 976
longevity, 800
Nabis roseipennis, 536
Pinus densiflora, 976
Pinus thunbergiana, 976
population dynamics, 976
predators, 976
Schizaphis graminum, 567
temperature, 800
Feeding, *Alabama argillacea*, 128
Blattella germanica, 198
Conopthorus resinosae, 88
Feeding, cotton leaves, 128
Dioryctria disclusa, 88
Nabis roseipennis, 536
Nilaparvata lugens, 678
Orius insidiosus, 532
Otiorhynchus sulcatus, 1074

oviposition, 1074
Panonychus citri, 190
Panonychus ulmi, 190
Schizaphis graminum, 118
Scirtothrips citri, 763
Sitophilus zeamais, 654
Spodoptera frugiperda, 182
Tetranychus pacificus, 190
Tetranychus urticae, 190
Feeding deterrents, *Leptinotarsa decemlineata*, 1057
Solanum chacoense, 1057
Feeding rhythms, biology, 914
density-related stress, 914
Lymantria dispar, 914
larval development, 914
Feeding success, *Aedes atlanticus*, 874
Aedes vexans, 874
body size, 874
Culex salinarius, 874
Psorophora columbiae, 874
Feltia jaculifera, pheromone, 40
traps, 40
Filberts, chemical exclusion, 1037
Myrocallis coryli, 1037
predators, 1037
Flight, *Cyclocephala pasadenae*, 194
Dendroctonus terebrans, 710
Diabrotica virgifera virgifera, 620
Pectinophora gossypiella, 494
Pholetesor ornigis, 331
Phyllophaga crinita, 194
Flight patterns, *Cheumatopsyche pettiti*, 355
swarming, 355
3-D technique, 355
Fluazifop, *Epilachna varivestis*, 216
Foliar biomass, *Leptinotarsa decemlineata*, 726
Foliar nitrogen, *Leptinotarsa decemlineata*, 726
Food effects, on *Platynota idaeusalis*, 210
Food consumption, *Chilomenes sexmaculata*, 800
development period, 800
fecundity, 800
longevity, 800
temperature, 800
Foraging, radioactive labeling, 1279
Solenopsis invicta, 1279
Forest structure, *Boisea rubrolineatus*, 1055
population dynamics, 1055
Forests, *Ceratocystis*, 1090
disturbance, 1090
epidemiology, 1090
Frankliniella occidentalis, Cotton, 821
functional response, 821
host range, 1292
Leucaena glauca, 1292
omnivore, 821
opportunist, 821
predator, 821
spider mites, 821
tomato spotted wilt virus, 1292
Fraser fir, *Adelges piceae*, 911
water stress, 911
Functional response, *Frankliniella occidentalis*, 821
omnivore, 821
opportunist, 821
predator, 821
spider mites, 821
Fungi, biological control, 1070
Nilaparvata lugens, 1070
rice, 1070
Fungicides, *Apis mellifera*, 1047
apples, 1047
pears, 1047

Gardenias, *Dialeurodes citri*, 585
Encarsia lahorensis, 585
Geocoris punctipes, *Anticarsia gemmatalis*, 1032
host-plant resistance, 1032
Pseudoplusia includens, 1032
soybeans, 1032
predators, 1032
Germ plasm, insect-resistant, 1, 11
simulation model, 1,11
Glandular chemicals, *Alabama argillacea*, 128
Glypta fumiferanae, *Chonistoneura pinus*, 409
host, 481
parasitization, 481
spatial activity, 409, 481
temporal activity, 409, 481
traps, 481
Grain sorghum, *Elasmopalpus lignosellus*, 905
larval distribution, 905
larval phenology, 905
Lygus desertinus, 132
Lygus elisus, 132
Lygus hesperus, 132
Graminella nigrifrons, corn, 1119
johnsongrass, 1119
life tables, 1119
Grass resistance, to *Spodoptera frugiperda*, 182
Grasshoppers, phenology, 924
Gretchena bolliana, larval infestion, 436
on pecan cultivars, 436
Growth, *Oligonychus pratensis*, 79
Growth inhibitors, artificial seeds, 1286
Callosobruchus maculatus, 1286
cowpeas, 1286
Guayule, *Lygus desertinus*, 132
Lygus elisus, 132
Lygus hesperus, 132
Guilds, *Dendroctonus* spp., 850
disturbance, 850
Ips spp., 850
population dynamics, 850

Haematobia irritans, circadian pattern, 1307
cow color, 1307
diapause survival, 44
overwintering, 44
populations, 44
Harpalus pennsylvanicus, beans, 1028
Evarthrus sodalis, 1028
intercropping, 1028
tomatoes, 1028
Harrisina americana, *Acoloithus falsarius*, 959
attractant, 959
pheromone, 959
Heliothis phloxiphaga, behavior, 931
pheromone identification, 931
sex pheromone, 931
Heliothis spp., *Eucelatoria bryani*, 1111
parasitoids, 1111
proteins, 1111
Heliothis virescens, *Bacillus thuringiensis*, 1161
bioassay, 1161
diapause, 919
Heliothis zea, 919, 1161
host, 476
host acceptance, 274
Microplitis croceipes, 476
nuclear polyhedrosis virus, 1161
parasitization by *Microplitis croceipes*, 274
parasitization, 476
thermal effects, 919
variation in density, 476
Heliothis zea, aflatoxin, 1116
Aspergillus flavus, 1116
Bacillus thuringiensis, 1161
bioassay, 1161
cotton, 365
cotton trichome density, 365
diapause, 919
Heliothis virescens, 919, 1161
host, 476
maize husk, 1116
Microplitis croceipes, 476
mortality, 365
Nabis roseipennis, 536
nuclear polyhedrosis virus, 1161
Ostrinia nubilalis, 1116
parasitization, 365
Trichogramma pretiosum, 365
thermal effects, 919
variation in density, 476
Herbicide effects, *Neochetina bruchi*, 1192
Neochetina eichhorniae, 1192
Herbicides, *Epilachna varivestis*, 216
Hippodamia convergens, *Nezara viridula*, 739
Hippodamia quinquesignata, control by *Verticillium lecanii*, 380
Horned larks, *Agrotis orthogonia*, 147
brain cholinesterase activity, 147
Host preferences, of Hymenoptera, 64
Host insect/plant relationships, *Bracon mellitor*, 880
boll weevil, 880
Host location, chemo-orientation, 839
Diabrotica virgifera virgifera, 839
olfaction, 839
Host odors, aggregation, 1050
Hylobius abietis, 1050
Host parasitism, *Musca domestica*, 436
Nasonia vitripennis, 436
Host plants, *Dacus cucurbitae*, 488
Host preference, endophytic fungus, 1083
Spodoptera frugiperda, 1083
tall fescue, 1083
Host range, bindweed, 834
Frankliniella occidentalis, 1292
Leucaena glauca, 1292
Megacerus discoidus, 834
tomato spotted wilt virus, 1292
Host suitability, elm, 843
Xanthogaleruca luteola, 843
Host/parasite effects, *Artogeia rapae*, 1234
larval density, 1234
Host-plant phenology, *Anticarsia gemmatalis*, 784
color phase, 784
density, 784
larval development, 784
Host-plant resistance, *Anticarsia gemmatalis*, 1032
Geocoris punctipes, 1032
Pseudoplusia includens, 1032
predators, 1032
soybeans, 1032
House flies, biological control, 1296
Muscidifurax zaraptor, 1296
Spalangia cameroni, 1296
stable flies, 1296
Urolepis rufipes, 1296
Humidity, aphid eggs, 989
Brachycolus asparagi, 989
Brachycorynella asparagi, 989
Diabrotica undecimpunctata howardi, 626
Diabrotica virgifera virgifera, 626
desiccation, 989
Stomoxys calcitrans, 706
Verticillium lecanii, 380
Hylobius abietis, aggregation, 1050
host odors, 1050
Hymenoptera, colored sticky panels, 64
forest, 64
host preferences, 64
microhabitat preferences, 64
parasitic, 64
Hypera postica, alfalfa resistance, 396
Medicago disciformis, 396
Medicago glandulosa, 396
Medicago glutinosa, 396
Medicago prostrata, 396
Medicago sativa, 396

Medicago scutellata, 396
on glandular-haired *Medicago* spp., 396
Hyperparasitism, *Apanteles fumiferanae*, 662
Apanteles morrisi, 662

Induced susceptibility, light, 803
soybeans, 803
Trichoplusia ni, 803
Infection, *Verticillium lecanii*, 380
Insecticide resistance, biological control, 1255
pesticide-resistant natural enemies, 1255
phytoseiid mites, 1255
Insecticides, *Apis mellifera*, 170
methyl parathion, 170
Penncap-M, 170
toxicity, 170
Integrated pest management, *Musca domestica*, 56
stochastic model, 540
Intercropping, beans, 1028
Evarthrus sodalis, 1028
Harpalus pennsylvanicus, 1028
tomatoes, 1028
Interspecific competition, *Cycloss turbinata*, 177
Metagneira grinnelli, 177
Ips calligraphus, associates, 1104
exclusion, 1104
mortality, 1104
Ips spp., beneficial insects, 1220
Dendroctonus spp., 850
disturbance, 850
guilds, 850
pine trees, 1220
population dynamics, 850

Johnsongrass, corn, 1119
Graminella nigrifrons, 1119
life tables, 1119

Laplatacris dispar, density relationships, 775
mortality, 775
population dynamics, 775
Larval density, *Artogeia rapae*, 1234
host/parasite effects, 1234
Mayetiola destructor, 305
Larval development, *Anticarsia gemmatalis*, 784
biology, 914
color phase, 784
density, 784
density-related stress, 914
feeding rhythms, 914
host-plant phenology, 784
Lymantria dispar, 914
Larval distribution, *Elasmopalpus lignosellus*, 905
grain sorghum, 905
larval phenology, 905
Larval phenology, *Elasmopalpus lignosellus*, 905
grain sorghum, 905
larval distribution, 905
Lepidoptera, synchrononous population fluctuations, 443
Leptinotarsa decemlineata, colonization, 311
feeding deterrants, 1057
foliar biomass, 726
foliar nitrogen, 726
plant feeding, 894
plant spacing effects, 311
planting date effects, 311
Podisus maculiventris, 894
potato persistence, 726
potatoes, 726
predatory pentatomids, 894
Solanum chacoense, 1057
Leptopilina boulardi, *Drosophila melanogaster*, 859
Drosophila simulans, 859
seasonal development, 859
synchrony, 859
Leucaena glauca, *Frankliniella occidentalis*, 1292
host range, 1292
tomato spotted wilt virus, 1292
Life cycle, *Anaphes sordidatus*, 100
Life stage, *Scirtothrips citri*, 763
Life tables, corn, 1119
fecundity, 976
Graminella nigrifrons, 1119
johnsongrass, 1119
Pinus densiflora, 976
Pinus thunbergiana, 976
population dynamics, 976
predators, 976
Light, induced susceptibility, 803
soybeans, 803
Trichoplusia ni, 803
Light intensity, *Aonidiella aurantii*, 971
male emergence, 971
relative humidity, 971
temperature, 971
Ligyrus subtropicus, *Bacillus popilliae*, 703
Cyclocephala parallela, 703
sugarcane, 703
Lima beans, *Epilachna varivestis*, 216
Liriomyza trifolii, bell pepper infestation, 383
bell peppers, 96
binomial sampling plans, 268
dispersion pattern, 383
in chrysanthemums, 268
within-plant distribution, 96
Listronotus oregonensis, *Anaphes sordidatus*, 100
Anaphes sordidatus, 469
host, 469
mortality, 469
overwintering sites, 113
parasitization, 469
survival, 113

Longevity, *Chilomenes sexmaculata*, 800
development period, 800
Elasmopalpus lignosellus, 715
egg parasites, 1149
fecundity, 800
food consumption, 800
Lymantria dispar, 1149
Nabis roseipennis, 536
Ooencyrtus kuvanae, 1149
photoperiod, 1149
progeny production, 1149
Schizaphis graminum, 567, 756
temperature, 800
Low-intensity radiation, *Aphidoletes aphidimyza*, 1067
diapause, 1067
Lumbricus terrestris, leaf burial, 321
mortality of *Pholetesor ornigis*, 321
mortality of *Pholetesor pedias*, 321
mortality of *Phyllonorycter blancardella*, 321
Lupinus arcticus, *Macrosiphum albifrons*, 719
Lupinus polyphyllus, *Macrosiphum albifrons*, 719
Lupinus sericeus, *Macrosiphum albifrons*, 719
Lygus desertinus alfalfa, 132
cotton, 132
grain sorghum, 132
guayule, 132
parasites of, 132
Lygus elisus, alfalfa, 132
cotton, 132
grain sorghum, 132
guayule, 132
parasites of, 132
Lygus hesperus, alfalfa, 132
cotton, 132
grain sorghum, 132
guayule, 132
parasites of, 132
Lygus lineolaris, *Nabis roseipennis*, 536
Lymantria dispar, between-tree movement, 373
biology, 914
burlap bands, 373
density/height profile, 431
density-related stress, 914
dispersal, 431
egg parasites, 1149
feeding rhythms, 914
larval development, 914
longevity, 1149
Ooencyrtus kuvanae, 1149
photoperiod, 1149
population, 122
progeny production, 1149
sampling, 122

Macrosiphum albifrons, *Aphidius lupini*, 719
Entomophthora sp., 719
Ephedrus californicus, 719
Lupinus arcticus, 719
Lupinus polyphyllus, 719
Lupinus sericeus, 719
mortality, 719
Praon sp., 719
parasitization, 719
Macrosiphum avenae, *Rhopalosiphum maidis*, 954
Rhopalosiphum padi, 954
Schizaphis graminum, 954
sampling, 954
spatial distribution, 954
Maize, *Chilo partellus*, 1124
Eldana saccharina, 1124
resistance, 1124
Maize husk, aflatoxin, 1116
Aspergillus flavus, 1116
Heliothis zea, 1116
Catrinia nubilalis, 1116
Malacosoma americanum, energy budget, 795
nutritional ecology, 795
Malacosoma disstria, parasites, 1184
population fluctuation, 1184
Male behavior, Araneae, 1225
ballooning, 1225
dispersal, 1225
Male emergence, *Aonidiella aurantii*, 971
light intensity, 971
relative humidity, 971
temperature, 971
Management, *Miliarevata lugens*, 93
Manduca sexta, compensatory processes, 1137
Nicotiana tabacum, 1137
simulated defoliation, 1137
yield loss, 1137
Mayetiola destructor, effect on wheat cold-hardiness, 392
genetic model, 24
in wheat, 305
insect-resistant germ plasm, 11
larval density, 305
model, 11
on wheat sugar reserves, 392
resistance, 24, 305
Triticum aestivum, 392
temperature effects, 305
wheat, 11, 24
Mefluidide, *Epilachna varivestis*, 216
Meganerus discoidus, bindweed, 834
host range, 834
Melanoplus sanguinipes, control by *Verticillium lecanii*, 281
eggs, 205
parasitization, 205
Melanotus communis, movement, 462
radiographs, 462
Metarhizium anisopliae, *Beauveria bassiana*, 1237
Chrysomelidae, 1237
sunlight, 1237

Metepeira grinnelli, biological control, 177
Cyclosa turbinata, 177
interspecific competition, 177
on insect populations, 177
predators, 177
Methyl parathion, *Apis mellifera*, 170
Penncap-M, 170
toxicity, 170
Metopolophium dirhodum, control by *Verticillium lecanii*, 281
Metopolophium persicae, control by *Verticillium lecanii*, 281
Microhabitat preferences, of Hymenoptera, 64
Microhabitat distribution, *Chromaphis juglandicola*, 555
Microplitis croceipes, fitness components, 274
Heliothis virescens, 476
Heliothis zea, 476
host acceptance, 274
host preference, 274
host, 476
linear function response, 476
parasitization of *Heliothis virescens*, 274
parasitization, 476
variation in *Heliothis* spp. density, 476
Migration, adult diapause, 884
Beauveria bassiana, 884
boundary layer, 1199
Cicadellidae, 1199
Conotrachelus nenuphar, 884
oogenesis/flight syndrome, 1199
Miscogaster hortensis, *Agromyza frontella*, 448
biological control, 448
Mocis disseverans, *Anticarsia gemmatalis*, 1272
sex pheromone, 1272
Model, genetic, 24
integrated pest management, 540
Mayetiola destructor, 24
phenology, 540
stochastic, 540
Moisture, conidial germination, 1154
Delia antiqua, 1154
Entomophthora muscae, 1154
temperature, 1154
Monochamus carolinensis, development, 251
seasonal occurrence, 251
Monomorium minimum, *Nezara viridula*, 739
Monoterpene, *Dendroctonus terebrans*, 710
Mortality, *Apanteles fumiferanae*, 662
Apanteles morrisi, 662
Apis mellifera, 166, 170
associates, 1104
Blattella germanica, 221
density relationships, 775
Entomophthora sp., 719
Epilachna varivestis, 32
Eumegachile pugnata, 877
exclusion, 1104
Ips calligraphus, 1104
Lepidoptera diaspar, 775
Lumbricus terrestris, 321
Macrosiphum albifrons, 719
nesting, 877
Oryctes rhinoceros, 84
Pholetesor ornigis, 321
Pholetesor pedias, 321
Phyllonorycter blancardella, 321
population dynamics, 775
temperature, 877
Moths, synchrononous population fluctuations, 443
Musca autumnalis, diapause commitment, 669
diapause induction, 669
temperature, 669
Musca domestica, *Cholepis rufipes*, 590
diel activity, 56
early season populations, 590
integrated pest management, 56
Muscidifurax zaraptor, 590
Nasonia vitripennis, 436
parasitization, 590
poultry manure, 56
Spalangia cameroni, 590
Stomoxys calcitrans, 590
seasonal activity, 56
Muscidifurax zaraptor, biological control, 1296
cold acclimation, 936
diapause, 936
house flies, 1296
Musca domestica, 590
Spalangia cameroni, 936, 1296
stable flies, 1296
Urolepis rufipes, 936, 1296
Myzocallis coryli, chemical exclusion, 1037
filberts, 1037
predators, 1037
Myzus persicae, abundance, 49
nitrogen effects, 49
population, 49
potatoes, 49
Verticillium lecanii, 380
within-plant distribution, 49

Nabis alternatus, control by *Verticillium lecanii*, 281
Nabis roseipennis, development, 536
diet, 536
fecundity, 536
feeding, 536
Heliothis zea, 536
longevity, 536
Lygus lineolaris, 536
temperature, 536
Nasonia vitripennis, biological control, 436
host parasitism, 436
Musca domestica, 436
progeny production, 436
Neochetina bruchi, herbicide effects, 1192
Neochetina eichhorniae, 1192

Neochetina eichhorniae, herbicide effects, 1192
Neochetina bruchi, 1192
Nephotettix nigropictus, parasitism in rice, 263
seasonal abundance, 263
Nephotettix virescens, development, 417
on modern rice cultivars, 417
on traditional rice cultivars, 417
population, 417
Nesting, *Eumegachile pugnata*, 877
mortality, 877
temperature, 877
Nezara viridula, aggregation, 739
development, 739
Hippodamia convergens, 739
Monomorium minimum, 739
Pheidole dentatum, 739
Podisus maculiventris, 739
Solenopsis invicta, 739
survival, 739
Tertamorium guineense, 739
Nicotiana tabacum, compensatory processes, 1137
Manduca sexta, 1137
simulated defoliation, 1137
yield loss, 1137
Nilaparvata lugens, biological control, 1070
development, 417
feeding, 678
fungi, 1070
management, 93
on modern rice cultivars, 417
on traditional rice cultivars, 417
parasitism in rice, 263
population, 93, 417, 648
rice, 648, 678, 1070
rice cultivars, 93
rice resistance, 648
seasonal abundance, 263
Nitrogen effects, on *Myzus persicae*, 49
Nosema epilachna, *Epilachna varivestis*, 32
Nosema varivestis, 32
Pediobius foveolatus, 32
Nosema varivestis, *Epilachna varivestis*, 32
Nosema epilachnae, 32
Pediobius foveolatis, 32
Nuclear polyhedrosis virus, *Anticarsia gemmatalis*, 573
Heliothis virescens, 1161
Spodoptera exigua, 240
soil-borne, 573
soybeans, 573
Nutritional ecology, energy budget, 795
Malacosoma americanum, 795
Nymphula depunctalis, development, 417
on modern rice cultivars, 417
on traditional rice cultivars, 417
population, 417

Oats, *Thrips tabaci*, 695
Olfaction, chemo-orientation, 839
Diabrotica virgifera virgifera
host location, 839
Oligonychus pratensis, corn, 79
density, 79
dispersion patterns, 335
field corn, 335
growth, 79, 335
plant stress, 79
sampling, 335
temperature, 79
Omnivore, cotton, 821
Franklinielle occidentalis, 821
functional response, 821
opportunist, 821
predator, 821
spider mites, 821
Ooencyrtus kuvanae, egg parasites, 1149
Lymantria dispar, 1149
longevity, 1149
photoperiod, 1149
progeny production, 1149
Oogenesis/flight syndrome, boundary layer, 1199
Cicadellidae, 1199
migration, 1199
Operophtera brumata, *Agrypon flaveolatum*, 1042
biological control, 1042
Cyrenis albicans, 1042
parasitism, 1042
Opportunist, cotton, 821
Franklinielle occidentalis, 821
functional response, 821
omnivore, 821
predator, 821
spider mites, 821
Orius insidiosus, apples, 984
developmental thresholds, 984
feeding, 532
functional response, 532
Panonychus ulmi, 532
predation, 984
temperature effects, 532
Oryctes rhinoceros, abundance, 84
breeding, 84
coconuts, 84
mortality, 84
survival, 84
Ostrinia nubilalis, aflatoxin, 1116
Aspergillus flavus, 1116
control by *Verticillium lecanii*, 281
corn, 106, 501, 631, 817
damage on cotton, 399
distribution, 399
egg masses, 501
egg-mass detrition, 817
egg-mass dispersion, 1212
feeding behavior, 399

Heliothis zea, 1116
larvae, 501
maize rusk, 1116
oviposition preference, 631
population, 501
sampling, 1212
sunflower, 631
survival, 631
within-field distribution, 106
within-plant distribution, 106
Otiorhynchus sulcatus, feeding, 1074
oviposition, 1074
Ovarian parameters, *Aphis pomi*, 607, 614
Overfeeding, arrested larval development, 1264
crowding, 1264
Tribolium castaneum, 1264
Overwintering, *Rivellia quadrifasciata*, 349
Thrips tabaci, 695
Trichogramma exiguum, 659
Overwintering sites, *Listronotus oregonensis*, 113
Oviposition, *Bruchophagus roddi*, 388
Elasmopalpus lignosellus, 715
feeding, 1074
Ostrinia nubilalis, 631
Otiorhynchus sulcatus, 1074
sites for *Eoreuma loftini*, 75
Sitona lineatus, 601
Spodoptera frugiperda, 327
Thrips tabaci, 695
Oviposition behavior, chrysanthemums, 1189
Spodoptera exigua, 1189

Panonychus citri, almond leaves, 190
damage, 190
feeding, 190
Panonychus ulmi, almond leaves, 190
damage, 190
feeding, 190
Orius insidiosus, 532
prey preference, 967
Stethorus punctum, 967
Tetranychus urticae, 967
temperature effects, 532
Papaipema nebris, diapause development, 403
eggs, 403
postdiapause development, 403
Paraquat, *Dendroctonus terebrans*, 710
Parasites, *Bemisia tabaci*, 1179
biological control, 1268
developmental rate, 1179
Malacosoma disstria, 1184
plant viruses, 1179
population fluctuation, 1184
Rhopalomyia californica, 1268
Tetrastichus sp., 1268
Torymus baccharidis, 1268
weed hosts, 1179
Parasitism, *Agrypon flaveolatum*, 1042
Artogeia rapae, 994
biological control, 1042
by *Bessa harveyi*, 246
by *Nephotettix nigropictus*, 263
by *Nilaparvata lugens*, 263
by *Pholetesor ornigis*, 331
by *Sogatella furcifera*, 263
Cyzenis albicans, 1042
in rice, 263
Operophtera brumata, 1042
of *Phyllonorycter blancardella*, 331
of *Pikonema alaskensis*, 246
Phyllonorycter elmaella, 635
Psigalio flavipes, 635
Pteromalus puparum, 994
Sympiesis marylandensis, 635
Parasitization, *Anaphes sordidatus*, 469
Apanteles fumiferanae, 662
Apanteles morrisi, 662
Aphidius lupini, 719
Ephedrus californicus, 719
Glypta fumiferanae, 481
Heliothis virescens, 476
Heliothis zea, 476
Listronotus oregonensis, 469
Macrosiphum albifrons, 719
Melanoplus sanguinipes, 205
Microplitis croceipes, 476
Musca domestica, 590
Muscidifurax zaraptor, 590
Praon sp., 719
Spalangia cameroni, 590
Spodoptera frugiperda, 342
Stomoxys calcitrans, 590
Urolepis rufipes, 590
Parasitoids, aphids, 1171
Eucelatoria bryani, 1111
Heliothis spp., 1111
proteins, 1111
Peanuts, *Bracon hebetor*, 767
Cadra cautella, 767
Plodia interpunctella, 767
Xylocoris flavipes, 767
Pear oviposition, *Psylla pyricola*, 1109
Pears, *Apis mellifera*, 1047
fungicides, 1047
Pecans, *Gretchena bolliana*, 436
Pectinophora gossypiella, adult flight phenology, 494
diapause, 494
termination, 494
Penncap-M, Apis mellifera, 170
toxicity, 170
Perkinsiella saccharicida, population dynamics, 1100
temperature effects, 1100
Pest management, *Leptinotarsa decemlineata*, 311
Stibadium spumosum, 1275
sunflowers, 1275
Pesticide-resistant natural enemies, biological control, 1255
insecticide resistance, 1255
phytoseiid mites, 1255
Pesticides, *Chrysoperla carnea*, 1130
carbaryl, 1130
resistance, 1130
Pheldole dentatum, *Nezara viridula*, 739
Phenology, *Diabrotica virgifera virgifera*, 109
grasshoppers, 924
stochastic model, 540
Phenology models, *Acyrthosiphon kondoi*, 227
Acyrthosiphon pisum, 227
Pheromone, *Acoloithus falsarius*, 959
Alsophila pometaria, 943
attractant, 959
behavior, 943
Choristoneura fumiferana, 152
communication disruption, 943
Feltia jaculifera, 40
Harrisina americana, 959
traps, 40
Pheromone identification, behavior, 931
Heliothis phloxiphaga, 931
sex pheromone, 931
Pheromones, *Agyrotaenia citrana*, 232
Dioryctria amatella, 316
Sitophilus zeamais, 733
Pholetesor ornigis, flight, 331
Lumbricus terrestris effects on mortality, 321
parasitoid of *Phyllonorycter blancardella*, 331
trapping, 331
Pholetesor pedias, *Lumbricus terrestris* effects on mortality, 321
Photoperiod, egg parasites, 1149
Lymantria dispar, 1149
longevity, 1149
Ooencyrtus kuvanae, 1149
progeny production, 1149
Phyllonorycter blancardella, *Lumbricus terrestris* effects on mortality, 321
parasitized by *Pholetesor ornigis*, 331
Phyllonorycter crataegella, monitoring, 562
population, 562
visual traps, 562
Phyllonorycter elmaella, apple, 635
Psigalio flavipes, 635
parasitism, 635
Symplesis marylandensis, 635
Phyllophaga crinita, Cyclocephala pasadenae, 194
flight, 194
traps, 194
Phyllotreta cruciferae, on cabbage in bare-ground monocultures, 293
on cabbage in living mulches, 293
Physiological time, computer program, 999
degree-day, 999
thermal unit, 999
Phytoseiid mites, biological control, 1255
insecticide resistance, 1255
pesticide-resistant natural enemies, 1255
Phytoseilus persimilis, *Amblyseius californicus*, 790
biological control, 790
corn, 790
Oligonychus pratensis, 790
Tetranychus urticae, 790
Pieris rapae, on cabbage in living mulches, 293
Pikonema alaskensis, life table, 246
parasitization by Bessa harveyi, 246
Pine, *Dioryctria amatella*, 316
Pine trees, beneficial insects, 1220
Ips spp., 1220
Pinus densiflora, fecundity, 976
life tables, 976
Pinus thunbergiana, 976
population dynamics, 976
predators, 976
Pinus thunbergiana, fecundity, 976
life tables, 976
Pinus densiflora, 976
population dynamics, 976
predators, 976
Plant feeding, *Leptinotarsa decemlineata*, 894
Podisus maculiventris, 894
predatory pentatomids, 894
Plant resistance, *Schizaphis graminum*, 581
Plant stress, *Oligonychus pratensis*, 79
Plant susceptibility, aphids, 1145
discriminant analysis, 1145
shade trees, 1145
urban entomology, 1145
Plant viruses, *Bemisia tabaci*, 1179
developmental rate, 1179
parasites, 1179
weed hosts, 1179
Platynota idaeusalis, food effects, 210
nondiapause larval development, 210
nondiapause pupal development, 210
sex effects, 210
strain effects, 210
thermal requirements, 210
Plodia interpunctella, *Bracon hebetor*, 767
male response, 369
natural enemies, 767
peanuts, 767
pheromone communication, 369
temporal differences, 369
wind tunnel, 369
Xylocoris flavipes, 767
Psigalio flavipes, *Phyllonorycter elmaella*, 635
Podisus maculiventris, *Leptinotarsa decemlineata*, 894
Nezara viridula, 739
plant feeding, 894

Podisus maculiventris, predatory pentatomids, 894
Pollen transfer, *Apis mellifera*, 723
Polyethylene glycol, *Schizaphis graminum*, 756
Polyhedrosis virus, *Bacillus thuringiensis*, 1161
  bioassay, 1161
  Heliothis zea, 1161
Polymorphism, *Acleris minuta*, 258
Population, *Adelges cooleyi*, 254
  Agrotis orthogonia, 147
  Anasa tristis, 285
  Bemisia tabaci, 61
  Brevicoryne brassicae on cabbage, 293
  Ceratitis capitata, 507
  Chilo suppressalis, 417
  Choristoneura fumiferana, 152
  Cnaphalocrocis medinalis, 417
  Cyclosa turbinata, 177
  Diabrotica vergifera vergifera, 109
  Dialeurodes citri, 585
  Diaphania hyalinata, 285
  Haematobia irritans, 44
  Lepidoptera, 443
  Lymantria dispar, 122
  Metepeira grinnelli, 177
  moths, 443
  Mephotettix viriscens, 417
  Nilaparvata lugens, 417, 648
  Nilaparvata lugens, 93
  Nymphula depunctalis, 417
  Ostrinia nubilalis, 501
  Phyllonorycter crataegella, 562
  Phyllotreta cruciferae on cabbage, 293
  Pieris rapae on cabbage, 293
  Sogatella furcifera, 417
  synchronous fluctuations, 443
  Thrips tabaci, 513, 520
  Xanthogaleruca luteola, 143
Population dynamics, *Anticarsia gemmatalis*, 809
  Boisea rubrolineatus, 1055
  Dendroctonus spp., 850
  density relationships, 775
  disturbance, 850
  Empoasca fabae, 898
  fecundity, 976
  forest structure, 1055
  guilds, 850
  Ips spp., 850
  Lepletocris dispar, 775
  life tables, 976
  mortality, 775
  Perkinsiella saccharicida, 1100
  Pinus densiflora, 976
  Pinus thunbergiana, 976
  predators, 976
  temperature effects, 1100
Population effects, of cabbage in bare-ground monocultures, 293
  of cabbage in living mulches, 293
Population fluctuation, *Malacosoma disstria*, 1184
  parasites, 1184
Potato persistence, *Leptinotarsa decemlineata*, 726
Potatoes, *Leptinotarsa decemlineata*, 726
  Myzus persicae, 49
  nitrogen, 49
Poultry, *Musca domestica*, 56
Praon sp., *Macrosiphum albifrons*, 719
Predation, apples, 984
  carabids, 1017
  corn agroecosystem, 1017
  developmental thresholds, 984
  Orius insidiosus, 984
  predator activity, 1017
Predator, cotton, 821
  Frankliniella occidentalis, 821
  functional response, 821
  omnivore, 821
  opportunist, 821
  spider mites, 821
Predator activity, carabids, 1017
  corn agroecosystem, 1017
  predation, 1017
Predators, *Anticarsia gemmatalis*, 1032
  Cyclosa turbinata, 177
  chemical exclusion, 1037
  fecundity, 976
  filberts, 1037
  Geocoris punctipes, 1032
  host-plant resistance, 1032
  life tables, 976
  Metepeira grinnelli, 177
  Mysocallis coryli, 1037
  Pinus densiflora, 976
  Pinus thunbergiana, 976
  population dynamics, 976
  Pseudoplusia includens, 1032
  soybeans, 1032
Predatory pentatomids, *Leptinotarsa decemlineata*, 894
  Podisus maculiventris, 894
  plant feeding, 894
Prey preference, *Panonychus ulmi*, 967
  Stethorus punctum, 967
  Tetranychus urticae, 967
Progeny production, egg parasites, 1149
  Lymantria dispar, 1149
  longevity, 1149
  Ooencyrtus kuwanae, 1149
  photoperiod, 1149
Proteins, *Eucelatoria bryani*, 1111
  Heliothis spp., 1111
  parasitoids, 1111
Pseudoplusia includens, *Anticarsia gemmatalis*, 1032
  Geocoris punctipes, 1032
  host-plant resistance, 1032
  predators, 1032
  soybeans, 1032
Psorophora columbiae, *Aedes atlanticus*, 874
  Aedes vexans, 874
  body size, 874
  Culex salinarius, 874
  feeding success, 874
Psorophora ferox, breeding area, 683
  distribution, 683
  eggs, 683
  endaphic parameters, 683
Psylla pyricola, pear oviposition, 1109
Pteromalus puparum, *Artogeia rapae*, 994
  parasitism, 994
Pupation, *Atiagenus unicolor*, 1166
  survival, 1166

Rachiplusia ou, *Agrotis ipsilon*, 360
  Bonnetia comta, 360
  nuclear polyhedrosis virus, 360
Radiation, *Stomoxys calcitrans*, 706
Radioactive labeling, foraging, 1279
  Solenopsis invicta, 1279
Radiographs, *Melanotus communis*, 462
  movement analysis, 462
Rearing, parental *Aphis pomi*, 614
Relative humidity, *Aonidiella aurantii*, 971
  light intensity, 971
  male emergence, 971
  temperature, 971
Reproduction, *Aphis pomi*, 614
  Bombus flavifrons, 750
  Schizaphis graminum, 567, 756
Resistance, *Chilo partellus*, 1124
  carbaryl, 1130
  Chrysoperla carnea, 1130
  Eldana saccharina, 1124
  Hypera postica, 396
  in wheat, 305
  maize, 1124
  Mayetiola destructor, 305
  Nilaparvata lugens, 648
  pesticides, 1130
  rice, 648
Response, *Sitophilus zeamais*, 733
Rhopalomyia californica, biological control, 1268
  parasites, 1268
  Tetrastichus sp., 1268
  Torymus baccharidis, 1268
Rhopalosiphum maidis, *Macrosiphum avenae*, 954
  Rhopalosiphum padi, 954
  Schizaphis graminum, 954
  sampling, 954
  spatial distribution, 954
Rhopalosiphum nymphaeae, temperature effects, 1096
Rhopalosiphum padi, *Macrosiphum avenae*, 954
  Rhopalosiphum maidis, 954
  Schizaphis graminum, 954
  sampling, 954
  spatial distribution, 954
  control by *Verticillium lecanii*, 281
Rice, biological control, 1070
  fungi, 1070
  Nilaparvata lugens, 93, 648, 1070
  resistant to *Nilaparvata lugens*, 678
  susceptible to *Nilaparvata lugens*, 678
Rivellia quadrifasciata, overwintering, 349
  seasonal abundance, 349

Sampling, *Adelges cooleyi*, 254
  egg-mass dispersion, 1212
  Lymantria dispar, 122
  Macrosiphum avenae, 954
  Oligonychus pratensis, 335
  Ostrinia nubilalis, 1212
  Rhopalosiphum maidis, 954
  Rhopalosiphum padi, 954
  Schizaphis graminum, 954
  spatial distribution, 954
  Tetranychus urticae, 335
Scelionidae, seasonal abundance, 1250
  soybeans, 1250
Schizaphis graminum, antibiosis tests, 581
  biotype B, 700
  biotypic status, 567
  drought stress, 118, 756
  fecundity, 567
  feeding, 118
  growth chambers, 581
  'Largo' wheat, 700
  longevity, 567, 756
  Macrosiphum avenae, 954
  plant resistance, 581
  polyethylene glycol, 756
  Rhopalosiphum maidis, 954
  Rhopalosiphum padi, 954
  reproduction, 756
  reproductive maturity, 567
  resistance, 700
  sampling, 954
  spatial distribution, 954
  wheat, 118, 756
Scirtothrips citri, feeding, 763
  life stage, 763
  temperature, 763
Seasonal abundance, *Adelges cooleyi*, 254
  Scelionidae, 1250
  soybeans, 1250
Seasonal activity, *Musca domestica*, 56
Seasonal development, *Leptopilina boulardi* 859
  Drosophila melanogaster 859
  Drosophila simulans 859
  synchrony, 859
Sethoxydim, *Epilachna varivestis*, 216

Sewage sludge, arthropods, 826
  collards, 826
Sex effects, on *Platynota idaeusalis*, 210
Sex pheromone, *Anticarsia gemmatalis*, 1272
  behavior, 931
  Cochylis hospes, 1063
  Heliothis phloxiphaga, 931
  Mocis disseverans, 1272
  pheromone identification, 931
Shade trees, aphids, 1145
  discriminant analysis, 1145
  plant susceptibility, 1145
  urban entomology, 1145
Simulated defoliation, compensatory processes, 1137
  Manduca sexta, 1137
  Nicotiana tabacum, 1137
  yield loss, 1137
Simulation model, diploid, 1
  insect-resistant germ plasm, 1
  two-locus, 1
Site selection, *Bruchophagus roddi*, 388
Sitona lineatus, egg melanization rate, 601
  ovipositional rhythms, 601
Sitophilus zeamais, age, 733
  corn, 654
  crowding, 733
  dosage, 733
  feeding, 654
  pheromones, 733
  response, 733
  starvation, 733
  3-pentanone, 733
Sogatella furcifera, development, 417
  on modern rice cultivars, 417
  on traditional rice cultivars, 417
  parasitism in rice, 263
  population, 417
  seasonal abundance, 263
Solanum chacoense, feeding deterrents, 1057
  Leptinotarsa decemlineata, 1057
Solenopsis invicta, foraging, 1279
  Nezara viridula, 739
  radioactive labeling, 1279
Soybeans, *Anticarsia gemmatalis*, 1032
  Anticarsia gemmatalis, 573, 779, 809
  density, 779
  Epilachna varivestis, 216
  Geocoris punctipes, 1032
  host-plant resistance, 1032
  induced susceptibility, 803
  light, 803
  nuclear polyhedrosis virus, 573
  Pseudoplusia includens, 1032
  population dynamics, 809
  predators, 1032
  Scelionidae, 1250
  seasonal abundance, 1250
  Trichoplusia ni, 803
Spalangia cameroni, biological control, 1296
  cold acclimation, 936
  diapause, 936
  house flies, 1296
  Musca domestica, 590
  Muscidifurax zaraptor, 936, 1296
  stable flies, 1296
  Urolepis rufipes, 936, 1296
Spatial, *Glypta fumiferanae*, 481
Spatial abundance, *Adelges cooleyi*, 254
Spatial activity, *Apanteles fumiferanae*, 409
  Apanteles morrisi, 409
  Glypta fumiferanae, 409
Spatial dispersion, *Chromaphis juglandicola*, 555
Spatial distribution, *Macrosiphum avenae*, 954
  Rhopalosiphum maidis, 954
  Rhopalosiphum padi, 954
  Schizaphis graminum, 954
  sampling, 954
Spider mites, almond leaves, 594
  cotton, 821
  Franklinella occidentalis, 821
  functional response, 821
  gas-exchange rates of almond leaves, 594
  omnivore, 821
  predator, 821
  water potential of almond leaves, 594
  water stress, 594
Spiders, abundance, 527
  diversity, 527
  sugarcane, 527
  weed control practices, 527
Spodoptera exigua, behavior, 1189
  chrysanthemums, 1189
  nuclear polyhedrosis virus, 240
  oviposition behavior, 1189
Spodoptera frugiperda, *Archytas marmoratus*, 342
  artificial diet, 327
  bermudagrass, 889
  Campoletis sonorensis, 342
  Chelonus insularis, 342
  Cotesia marginiventris, 342
  corn, 342
  corn leaves, 327
  development, 182, 889
  endophytic fungus, 1083
  feeding, 182
  forage crops, 889
  grass resistance, 182
  host grasses, 182
  host preference, 1083
  larvae, 327
  larval grass, 327
  Ophion flavidus, 342
  overwintering areas, 342
  oviposition, 327
  parasitization, 342
  parasitoids, 342
  Rogas laphygmae, 342
  sorghum, 342
  spring distribution, 342
  Temelucha difficilis, 342
  tall fescue, 1083
Sporulation, *Verticillium lecanii*, 380
Squash, *Anasa tristis*, 285
  Diaphania hyalinata, 285
Stable flies, biological control, 1296
  house flies, 1296
  Muscidifurax zaraptor, 1296
  Spalangia cameroni, 1296
  Urolepis rufipes, 1296
Starvation *Sitophilus zeamais*, 733
Steinernema feltiae, *Ceratitis capitata*, 465
  Dacus cucurbitae, 465
  Dacus dorsalis, 465
Stethorus punctum, *Panonychus ulmi*, 967
  prey preference, 967
  Tetranychus urticae, 967
Stibadium spumosum, pest management, 1275
  sunflowers, 1275
Stochastic phenology model, *Choristoneura occidentalis*, 547
Stomoxys calcitrans, humidity effects, 706
  radiation effects, 706
  temperature effects, 706
  traps, 706
Strain effects, on *Platynota idaeusalis*, 210
Sugarcane, *Bacillus popilliae*, 703
  Cyclocephala parallela, 703
  Koreums loftini, 75
  Ligyrus subtropicus, 703
  spiders, 527
  weed control practices, 527
Sunflowers, *Ostrinia nubilalis*, 631
  pest management, 1275
  Stibadium spumosum, 1275
Sunlight, *Beauveria bassiana*, 1237
  Chrysomelidae, 1237
  Metarhizium anisopliae, 1237
Survival, *Alabama argillacea*, 128
  Agrypnus variicolor, 1166
  Diabrotica undecimpunctata howardi, 626
  Diabrotica virgifera virgifera, 626
  Listronotus oregonensis, 113
  Nezara viridula, 739
  Oryctes rhinoceros, 84
  Ostrinia nubilalis, 631
  pupation, 1166
Survivorship, biological control, 1301
  development rate, 1301
  Urolepis rufipes, 1301
Sweet corn, *Chaetocnema pulicaria*, 867
  traps, 867
Sympiesis marylandensis, *Phyllonorycter elmaella*, 635
Synchrony, *Leptopilina boulardi* 859
  rosophila melanogaster 859
  rosophila simulans 859
  seasonal development, 859

Tabanus nigrovittatus, visual attraction, 689
Tall fescue, endophytic fungus, 1083
  host preference, 1083
  Spodoptera frugiperda, 1083
Tarsonemus ips, bluestain, 951
  Ceratocystis minor, 951
  Dendroctonus frontalis, 951
  Tarsonemus krantzi, 951
Tarsonemus krantzi, bluestain, 951
  Ceratocystis minor, 951
  Dendroctonus frontalis, 951
  Tarsonemus ips, 951
Temperature, *Aonidiella aurantii*, 971
  Aphis pomi, 607, 614
  Bombus flavifrons, 750
  Chilomenes sexmaculata, 800
  conidial germination, 1154
  Delia antiqua, 1154
  Diabrotica undecimpunctata howardi, 626
  Diabrotica virgifera virgifera, 626
  development period, 800
  Elasmopalpus lignosellus, 715
  Entomophthora muscae, 1154
  Rumegachile pugnata, 877
  effects of food, 210
  effects of sex, 210
  effects of strain, 210
  fecundity, 800
  food consumption, 800
  light intensity, 971
  longevity, 800
  male emergence, 971
  Mayetiola destructor, 305
  moisture, 1154
  mortality, 877
  Musca autumnalis, 669
  Nabis roseipennis, 536
  nesting, 877
  Oligonychus pratensis, 79
  Orius insidiosus, 532
  Pancaychus ulmi, 532
  Platynota idaeusalis, 210
  relative humidity, 971
  Scirtothrips citri, 763
  Stomoxys calcitrans, 706
  Trichogramma exiguum, 659
Temperature effects, *Perkinsiella saccharicida*, 1100
  population dynamics, 1100
  Rhopalosiphum nymphaeae, 1096
Temporal, *Glypta fumiferanae*, 481
Temporal activity, *Apanteles fumiferanae*, 409
  Apanteles morrisi, 409
  Glypta fumiferanae, 409

Tertamorium guineense, *Nezara viridula*, 739
Tetranychus pacificus, almond leaves, 190
  damage, 190
  feeding, 190
Tetranychus urticae, almond leaves, 190
  damage, 190
  dispersion patterns, 335
  feeding, 190
  field corn, 335
  growth, 335
  Panonychus ulmi, 967
  prey preference, 967
  sampling, 335
  Stethorus punctum, 967
Tetrastichus sp., biological control, 1268
  parasites, 1268
  Rhopalomyia californica, 1268
  Torymus baccharidis, 1268
Theriaophis maculata, control by *Verticillium lecanii*, 281
Thermal unit, computer program, 999
  degree-day, 999
  physiological time, 999
Thrips tabaci, alfalfa, 513, 695
  cabbage, 513, 520
  ecology, 520
  oats, 513, 695
  overwintering, 695
  oviposition, 695
  population, 513, 520
  red clover, 513
  species composition, 520
  temporal occurrence, 513
  traps, 513
  wheat, 513, 695
Thuringiensin, *Apis mellifera*, 166
  B exotoxin, 166
  Bacillus thuringiensis, 166
  toxicity, 166
Tomato spotted wilt virus, *Frankliniella occidentalis*, 1292
  host range, 1292
  Leucaena glauca, 1292
Tomatoes, *Kvarthrus sodalis*, 1028
  Harpalus pennsylvanicus, 1028
  intercropping, 1028
Torymus baccharidis, biological control, 1268
  parasites, 1268
  Rhopalomyia californica, 1268
  Tetrastichus sp., 1268
Trap, *Bemisia tabaci*, 300
  design, 300
  placement, 300
  Trialeurodes abutilonea, 300
Trapping, *Ceratitis capitata*, 71
  Pholetesor ornigis, 331
Traps, *Ceratitis capitata*, 507
  Chaetocnema pulicaria, 867
  Choristoneura fumiferana, 152
  Dioryctria amatella, 316
  Glypta fumiferanae, 481
  Phyllonorycter crataegella, 562
  Phyllophaga crinita, 194
  sweet corn, 867
  Thrips tabaci, 513
  visual monitoring, 562
Tree species, aphid host preference, 771
Trialeurodes abutilonea, Bemisia tabaci, 300
  in cotton, 300
  in lettuce, 300
  traps, 300
Tribolium castaneum, arrested larval development, 1264
  crowding, 1264
  overfeeding, 1264
Trichogramma exiguum, overwintering, 659
  temperature effects, 659
Trichogramma pretiosum, cotton trichome density, 365
  cotton, 365
  mortality, 365
  parasitization of Heliothis zea, 365
Trichogramma retorridum, parasite of *Eoreuma loftini*, 75
Trichoplusia ni, induced susceptibility, 803
  light, 803
  soybeans, 803
Trimedlure, *Ceratitis capitata*, 71
Triticum aestivum, cold-hardiness, 392
  Mayetiola destructor, 392
  sugar reserves, 392
Turpentine, *Dendroctonus terebrans*, 710

Urban entomology, aphids, 1145
  discriminant analysis, 1145
  plant susceptibility, 1145
  shade trees, 1145
Urolepis rufipes, biological control, 1296
  biological control, 1301
  cold acclimation, 936
  development rate, 1301
  diapause, 936
  house flies, 1296
  Musca domestica, 590
  Muscidifurax zaraptor, 936, 1296
  Spalangia cameroni, 936, 1296
  stable flies, 1296
  survivorship, 1301

Verticillium lecanii, biological control agent, 380
  control agent, 281
  humidity, 380
  infection, 380
  Myzus persicae, 380
  on aphid populations, 281
  sporulation, 380
Visual attraction, *Tabanus nigrovittatus*, 689
Volatile components, alfalfa seed, 1244
  behavior, 1244
  chalcids, 1244

Water stress, *Adelges picea*, 911
  Fraser fir, 911
Weather, cotton, 1078
  Crocidosema plebejana, 1078
Weed control practices, spiders, 527
Weed hosts, *Bemisia tabaci*, 1179
  developmental rate, 1179
  parasites, 1179
  plant viruses, 1179
Weight, *Aphis pomi*, 607, 614
Wheat, *Agrotis orthogonia*, 147
  genetic model, 24
  'Largo', 700
  Mayetiola destructor, 11, 24, 305
  resistance, 24, 700
  Schizaphis graminum, 118, 700, 756
  Thrips tabaci, 695

Xanthogaleruca luteola, elm, 843
  elm preference, 143
  host suitability, 843
  population, 143
Xylocoris flavipes, *Cadra cautella*, 767
  natural enemies, 767
  peanuts, 767
  Plodia interpunctella, 767

Yeast infection, *Blattella germanica*, 221
Yield loss, compensatory processes, 1137
  Manduca sexta, 1137
  Nicotiana tabacum, 1137
  simulated defoliation, 1137
3-pentatone *Sitophilus zeamais*, 733

CUMULATIVE AUTHOR INDEX

Environmental Entomology, Volume 15, 1986

Abrahamson, L.P., 152
Adams, R.G., 867
Agnello, A.M., 216
Aguda, R.M., 1070
Alfiler, A., 84
Ali, A.D., 527
Alikhan, M.A., 800
AliNiazee, M.T., 1037, 1255
Allan, S.A., 689
Allen, D.C., 152
Ampofo, J.K.O., 1124
Anazonwu, D.L., 779
Anderson, D.L., 703
Andow, D.A., 293
Apperson, C.S., 683
Aquino, G.B., 93
Arbogast, R.T., 767
Arceo, M.B., 93
Archbold, E.F., 221
Arthur, A.P., 1063
Arthur, F.H., 911
Atkinson, K.D., 221
Auger, M., 152

Bacheler, J.S., 399
Backman, C.B., 715
Baker, J.E., 1166
Baker, T.C., 369
Ballou, J.K., 1096
Barfield, C.S., 327
Barnes, M.M., 190, 594
Barrett, B.A., 635
Barry, D., 1116
Beckwith, R.C., 540, 547
Bell, M.R., 1161
Benedict, J.H., 365
Bergman, M.K., 109, 839
Bergstrom, G., 1050
Berisford, C.W., 316
Berry, I.L., 706
Bloomfield, P., 683
Boethel, D.J., 1250
Boiteau, G., 311
Botella, L.M., 1264
Boucias, D.G., 703
Bowers, M.A., 750
Bradley, J.R. Jr., 216, 399
Bridges, J.R., 951
Broce, A.B., 706
Brooks, W.M., 32
Brust, G.E., 1017
Burkholder, W.E., 733
Burton, G.W., 889
Butler, G.D. Jr., 61
Butt, B.A., 1109
Buttery, R.G., 388, 1244
Byrne, D.N., 300

Calvin, D.D., 817, 1212
Cameron, R.S., 316
Campbell, J.B., 1307
Cantelo, W.W., 1057
Carroll, D.P., 607, 614, 1171
Carruthers, R.I., 1154
Canter, T.D., 1096
Chandler, L.D., 96, 383
Chang, N.T., 182
Cherry, R.H., 703, 1100
Chiang, H.C., 631
Chiang, H.S., 803
Cho, J.J., 1292
Clay, K., 1083
Coats, S.A., 620
Cohen, M.B., 719
Collins, R.D., 100, 113, 469
Cone, W.W., 989
Corrigan, J.E., 321
Cossentine, J.E., 360
Coudriet, D.L., 1179
Couey, H.M., 71
Coulson, R.N., 850
Cox, T.S., 24
Croft, B.A., 232, 1255
Culliney, T.W., 826
Cunningham, R.T., 71

Dahlsten, D.L., 1145
Danielson, S.D., 396
Daoust, R.A., 642, 1237
De Sagun, S., 93
Deahl, K.L., 1057
DeBarr, G.L., 316
Debolt, J.W., 132
Degrandi-Hoffman, G., 723
Dennis, B., 540, 547
Despins, J.L., 106
DeWeese, L.R., 147
Dorschner, K.W., 118, 756
Dos Santos, A., 1028
Doss, R.P., 1074
Drapek, R.J., 409
Drea, J.J. Jr., 448

Eckenrode, C.J., 1022
Edelson, J.V., 963
Eggen, D.A., 152
Ehler, L.E., 1268
Eidmann, H.H., 1050
Eikenbary, R.D., 118, 756
Elkinton, J.S., 371, 914
Elliott, N.C., 409, 662, 954
Epstein, M.E., 443

Fatzinger, C.W., 710
Federici, B.A., 240
Fescemyer, H.W., 784
Fisher, J.R., 626
Flamm, R.O., 850
Flexner, J.L., 232
Flinn, P.W., 898
Foard, D.E., 1286
Foster, R.E., 501
Freytag, P.H., 1119
Fried, C.S., 439
Fulton, W.C., 839
Funderburk, J.E., 905

Gage, S.H., 1078
Gaines, T.P., 889
Gallagher, K.D., 494
Gallun, R.L., 305
Gelernter, W.D., 240
Gilkeson, L.A., 1067
Gillott, C., 943
Gilstrap, F.E., 96, 335, 383, 790
Gould, F., 1, 11, 349, 462
Goyer, R.A., 1184, 1220
Grafius, E., 100, 113, 469
Grafton-Cardwell, E.E., 1130
Graham, H.M., 132
Green, T.A., 562
Guthrie, W.D., 1116
Gutierrez, A.P., 494, 555
Guzman, D.R., 936, 1296

Haag, K.H., 1192
Habeck, D.H., 182
Hadam, J.J., 1255
Hain, F.P., 911
Hajek, A.E., 771, 1145
Hall, R.W., 143, 843
Hamilton, J.G., 1078
Hamm, J.J., 207
Hammond, A.M., 784
Hammond, A.M. Jr., 1083
Hansen, E.M., 1090
Hanula, J.L., 316
Hardy, T.N., 1083
Harper, A.M., 281
Harris, E.J., 488, 507
Harrison, F.P., 1137
Hart, E.R., 258
Hatchett, J.H., 24
Haynes, D.L., 662, 1154
Heath, R.R., 327, 959, 1272
Heinrichs, E.A., 3, 417, 422, 648, 678
Hendrickson, R.M. Jr., 448
Heraty, J.M., 321
Hertlein, M.B., 859
Herzog, D.C., 905
Higley, L.G., 999
Hill, S.B., 1067
Hoffman, C.J., 300
Holtzer, T.O., 79
Hoopingarner, R., 723
Hopper, K.R., 274, 476
Horsburgh, R.L., 532, 984
Horwith, B., 1028
Houck, M.A., 967
Hower, A.A., 898
Hoxie, R.P., 392
Hoy, M.A., 1130
Hoyt, S.C., 607, 614, 1171
Huang, N.C., 281
Hudson, W.G., 585

Inayatullah, C., 581, 700
Isman, M.B., 971

Jackson, C.G., 132
Jansson, R.K., 49, 726
Johnson, J.R., 75
Johnson, R.C., 118, 756
Johnson, S.J., 779
Jones, V.P., 190, 268
Jones, W.A., 1250
Jorgensen, C.D., 635
Jung-Tsung, W., 648

Kamm, J.A., 388, 1244
Keever, D.W., 767
Keller, M.A., 659
Kelley, R.S., 152
Kemp, W.P., 540, 547, 924
Khan, Z.R., 803
Kieckhefer, R.W., 954
Kimball, B.A., 61
Kimberling, D.N., 1042
Kindler, S.D., 567
King, E.G., 476
Kishaba, A.N., 1179
Klomparens, K., 723
Klun, J.A., 931
Knapp, M.C., 817, 1212
Knight, A.L., 232
Koethe, R.W., 349
Kok, L.T., 834, 994, 1234
Kolodny-Hirsch, D.M., 122, 1137
Kovalev, B., 316
Kryszan, J.J., 427
Kuhna, A.S., 392
Kulman, H.M., 246
Kunz, S.E., 44
Kwolek, W.F., 1116

Laing, J.W., 321
Lance, D.R., 914
Landolt, P.J., 959, 1272
Lanier, G.N., 152
Lasota, J.A., 254, 994, 1234
Lee, C.Y.L., 507
Legg, D.E., 631, 654
Leigh, T.F., 821
Leonhardt, B.A., 931
Leppla, N.C., 1272
Letourneau, D.K., 285
Levine, E., 403
Lewis, L.C., 360
Liebhold, A.M., 373
Liljesthrom, G.G., 775
Lillehoj, E.B., 1116
Lindegren, J.E., 465
Linit, M.J., 251
Lisk, D.J., 826
Lockwood, J.A., 739
Long, G.E., 355
Lopez, J.D., 365, 931
Lopez, J.D. Jr., 919
Los, L.M., 867
Lunden, J.D., 1047
Lutton, G.G., 380
Lynch, R.E., 182, 889, 905

Mackauer, M., 719
Mack, T.P., 715, 905
Mahr, D.L., 331
Mandeville, J.D., 56
Manglitz, G.R., 396
Martin, P.B., 342
Mason, C.E., 170
Mason, P.G., 1063
Matsumoto, K., 959
Mattson, W.J., 88
Mauney, J.R., 61
Mawby, W.D., 219
Mayer, D.F., 1047
McCaffrey, J.P., 532, 984
McCartney, D.A., 1017
McClure, M.S., 976
McEwen, L.C., 147
McGiffen, M.E. Jr., 884
McGuinness, N., 1028
McMillan, W.W., 1116
McNew, R.W., 756
Medina, E.B., 417
Medrano, F.G., 417, 648
Meinke, L.J., 880
Mensua, J.L., 1264
Merchant, M.C., 795
Merkle, O.G., 700
Messing, R.H., 1037
Meyer, J.A., 56
Meyer, J.R., 884
Meyer, W.L., 316
Meyerdirk, D.E., 1179
Miller, J.C., 1042
Miller, M.C., 1104
Miller, R.W., 232

Miller, W.E., 443
Milner, R.J., 380
Mishoe, J.W., 809
Mitchell, H.R., 327
Mitchell, W.C., 1292
Mizell, R.F., 436
Monson, W.G., 889
Montandon, R., 128
Moon, R.D., 669
Morrison, R.K., 365
Morse, J.G., 763
Moser, J.C., 951
Mullen, M.A., 767
Mullens, B.A., 56
Murdock, L.L., 1286
Mutchmor, J.A., 620

Nadgauda, D., 536
Nasci, R.S., 874
Nation, J.L., 710
Nelson, A.K., 706
Nettles, W.C. Jr., 1111
Nguyen, H.T., 581
Nguyen, R., 1100
Nicholson, A.G., 293
Nishida, T., 488
Nordenhem, H., 1050
Nordlander, G., 1050
Norman, J.M., 79
Norris, D.M., 803
North, R.C., 513, 520, 695
Nowierski, R.M., 555
Nuessly, G.S., 1279
Nyrop, J.P., 481, 501

O'Keeffe, L.E., 601
Olien, C.R., 392
Onsager, J.A., 924
Oosterwyk, A.S., 803
Orr, D.B., 1250
Ostlie, K.R., 999
Own, O.S., 32

Pair, S.D., 342
Palaniswamy, P., 40, 943
Parker, F.D., 877
Parrella, M.P., 268
Payne, T.L., 850
Pedigo, L.P., 999
Pena, N., 263
Penrose, R.L., 1042
Pereira, R.M., 642, 1237
Perfecto, I., 1028
Perkins, W.D., 1275
Perring, T.M., 79
Pershing, J.C., 251
Petersen, J.J., 590, 936, 1296
Phelan, P.L., 369
Pickett, C.H., 335, 790
Pimentel, D., 439, 826
Pitre, H.N., 536
Pizzamiglio, M.A., 494
Plagens, M.J., 1225
Pomeroy, M.A., 1286
Ponelelt, C.G., 654
Poston, F.L., 817, 1212
Prabhaker, N., 1179
Press, J.W., 767
Price, J.R., 880
Prokopy, R.J., 562
Pulley, P.E., 850
Puterka, G.J., 880

Raina, A.K., 931
Rapusas, H.R., 417
Ratanatham, S., 305
Raulston, J.R., 342
Read, M.R., 669
Reagan, T.E., 527
Rees, N.E., 205
Reierson, D.A., 221
Reling, D., 431
Ridgway, N.M., 331
Riley, M.A., 1220
Roberts, D.W., 1070
Roberts, J.E. Sr., 106
Rock, G.C., 210
Rodriguez, J.G., 654
Roelofs, W.L., 316
Rogers, C.E., 1275
Rogers, D.J., 1032
Rombach, M.C., 1070
Romine, C.L., 1161
Ruberson, J.R., 894
Ruggiero, M.A., 795
Russin, J.S., 1250
Rust, M.K., 221
Rutz, D.A., 1301
Ryan, J.D., 118, 756
Rykiel, E.J., 850

Sanchez, N.E., 775
Sanford, L.L., 1057
Santos, W.J.D., 494
Saunders, J.A., 1137
Savinelli, C.E., 399
Schenk, T.C., 631
Schiffhauer, D.E., 436
Schladweiler, P., 147
Schotzko, D.J., 601
Schowalter, T.D., 1055, 1090
Schreiber, E.T., 1307
Schultz, B., 1028
Schwalbe, C.P., 914
Seaman, A., 501
Sedlacek, J.D., 1119
Segers, J.C., 365
Seiler, G.J., 1275
Shade, R.E., 1286
Shanks, C.H. Jr., 1074
Sheehan, W., 456
Shelton, A.M., 501, 513, 520, 695
Shepard, B.M., 1070
Shepard, M., 263
Shetlar, D.J., 254
Shimanuki, H., 166, 207
Shinn, E.A., 355
Siegfried, B.D., 710
Silverman, J., 198
Simmons, G.A., 409, 481, 662
Sinden, S.L., 1057
Slosser, J.E., 880
Smilowitz, Z., 49, 726
Smith, J.D., 1184
Smith, J.W. Jr., 75
Smith, L., 1301
Smits, P.H., 1189
Solomon, D.L., 683
Sonnet, P.E., 959
Sorensen, E.L., 396
Sosa, O. Jr., 1100
Sparks, A.N., 342
Spiller, D.A., 177
Spomer, S.M., 567
Steck, W.F., 40
Sterling, W.L., 128, 1279
Stimac, J.L., 809
Stinner, B.R., 1017
Stipanovic, R.D., 128
Stoffolano, J.G. Jr., 689
Stone, J.D., 194
Story, R.N., 739
Strnad, S.P., 839
Stuart, C., 1109
Suggars, A.L., 683
Sullivan, M.J., 1032
Sumner, L.C., 756
Sutter, G.R., 427
Swier, S.R., 152

Takara, J.M., 488
Tauber, C.A., 894
Tauber, M.J., 894
Taylor, R.A.J., 431, 898
Teraguchi, S.E., 1199
Thomas, D.B., 44
Throne, J.E., 1022
Tilles, D.A., 1050
Tipping, P.W., 654
Tollefson, J.J., 620
Treacy, M.F., 365
Trichilo, P.J., 821
Tsai, J.H., 1096
Turpin, F.T., 109

Underhill, E.W., 40, 943, 1063

Vail, P.V., 465
Valencia, S.L., 93
Valovage, W.D., 246
Van Duyn, J.W., 216, 349
Van Leerdam, M.B., 75
Vandenberg, J.D., 166, 207
Vandermeer, J., 1028
Vega, C.R., 417, 422
Velusamy, R., 678
Viajante, V., 417
Villacorta, A., 494
Villani, M.G., 462
Vinson, S.B., 128
Vlak, J.M., 1189
Von Bretzel, P.R., 300
van de Vrie, M., 1189

Wagner, T.L., 850
Walgenbach, C.A., 733
Waliner, W.E., 373
Wang, R., 834
Wassgren, A.-B., 1050
Weatherby, J.C., 258
Webster, J.A., 581, 700
Weiss, M.E., 803
Welch, S.M., 817, 1212
Wellso, S.G., 392
Welter, S.C., 190
Weseloh, R.M., 64, 1149
Widstrom, N.W., 1116
Wien, H.D., 293
Wiesenborn, W.D., 763
Wilkerson, G.G., 809
Wilkinson, R.C., 710
Williams, A.L., 327
Williams, H.J., 128
Williams, M.L., 585
Willson, H.R., 293
Wiseman, B.R., 182, 889
Witcosky, J.J., 1090
Wong, J.W., 943
Wright, L.C., 989

Xingquan, K., 817, 1212

Yan, J.-Y., 971
Yeargan, K.V., 1119
Yearian, W.C., 573
Young, C.E., 843
Young, S.Y., 573
Youngman, R.R., 190, 594
Yousuf, M., 800
Yudin, L.S., 1292

Zelazny, B., 84
Zuber, M.S., 1116